

FORTIFICATION OF MILK AND MILK PRODUCTS

P. Sathya¹, Lejaniya¹, C. Srinivasan^{2*}, V. Madhupriya³ and S. Sasikumar⁴

¹MVSc Scholars, Dept of Dairy Science, ²MVSc Scholar, Dept of Animal Nutrition,
College of Veterinary and Animal Science, Mannuthy, Thrissur - 680651

³MVSc Scholar, Dept of Poultry Science, ⁴Dept of Veterinary Clinical Medicine,
Ethics & Jurisprudence, Veterinary College and Research Institute, Namakkal – 637 002

E-mail: dr.srinivasan1992@gmail.com (* *Corresponding Author*)

Abstract: Fortification of milk with micronutrients play a major role in relieving nutritional deficiency problems in humans. It can improve the palatability and sensory of the products. The studies prove that wide use of fortified milk products in the world improve the health condition of people in geographically specific areas. In this review article fortification of milk and milk products with addition of iron, iodine, vitamins, fish oil, probiotic, conjugated linoleic acid, casinate and fiber fortification discussed below.

Keywords: Fortification, Milk Products, Iron, Vitamins, Probiotic.

INTRODUCTION

Fortification defines as "the practice of deliberately increasing the content of an essential micronutrient, i.e. vitamins and minerals (including trace elements). Food is the only source of all nutrients like water, carbohydrates, protein, fat, minerals and vitamins, required for the growth and maintenance of humans. The micronutrients which are found in the food, we consume on a regular basis, are sometimes available in lower amounts or not present in them. Hence, people suffer from deficiency diseases of these nutrients. The WHO and FAO recognized that there are over 2 billion people worldwide, who suffer from a variety of micronutrient deficiencies. Statistically discovered that approximately 1 in 3 people worldwide were at risk for either iodine, vitamin A, or iron deficiency. For obtaining compensation of micronutrients in the foods, innovation technologies like food fortification or enrichment of food products used through which these micronutrients are compensated in the regular diet of the common people.

FORTIFICATION IN MILK AND MILK PRODUCTS

The micronutrients which are commonly used in the fortification of the food products such as milk and its products are:

- Vitamins and co-vitamins

- Essential minerals
- Essential fatty acids
- Essential amino acids
- Phytonutrients
- Enzymes

METHODS OF FOOD FORTIFICATION

- **Bio Fortification** (i.e. breeding crops to increase their nutritional value - conventional selective breeding and modern genetic modification)
- **Synthetic Biology** (i.e. addition of probiotic bacteria to foods)
- **Commercial and Industrial Fortification** i.e. flour, rice, oils (common cooking foods).
- **Home Fortification** (e.g. vitamin D drops)

NUTRIENT FOR FORTIFICATION

IRON FORTIFICATION

Iron fortification of milk or dairy products induces several bio physicochemical modifications with important consequences. Iron fortification of food is regarded as the most cost-effective method for reducing the prevalence of nutritional iron deficiency. Nutritional anemia remains very prevalent in developing countries, and iron fortification appears until recently to have had little impact.

IODINE FORTIFICATION

Iodine concentration of organic summer milk was significantly lower than the iodine concentration of organic winter milk. No significant relation between iodine content and milk fat percentage with regard to geographical sampling location. Whey cheese iodine concentration was significantly higher than iodine concentration in casein cheeses. Goiter can be regulated by use of iodine fortified dairy products.

VITAMIN FORTIFICATION

Natural and added forms of vitamin A (all- trans-retinol, retinyl esters and beta-carotene) and vitamin E (alpha-tocopherol, alpha tocopheryl acetate) were determined in commercially available dairy products that are frequently consumed. Retinyl esters, beta carotene and alpha-tocopherol are in natural dairy products, whereas retinyl and alpha-tocopheryl acetate occur in most vitamin-fortified products.

Since Vitamin D is a fat-soluble vitamin, it cannot be added to a wide variety of foods. Foods that it is commonly added to margarine, vegetable oils and dairy products.

Rickets could be cured by these foods. Their results showed that sunlight exposure and cod liver oil were the cure. It was not until the 1930s that vitamin D was actually linked to curing rickets. This discovery led to the fortification of common foods such as milk, margarine, and breakfast cereals. The astonishing statistics of approximately 80–90% of children showing varying degrees of bone deformations due to vitamin D deficiency to being a very rare condition. The current RDA for infants aged 0–6 months is 10 μg (400 International Units (IU))/day and for adults over 19 years of age it is 15 μg (600 IU)/day.

FISH OIL FORTIFICATION

Fortification of dairy products with long-chain polyunsaturated fatty acid, however, the level of fortification was limited. The highest level of fortification was obtained for solid, high-fat dairy products (spreadable fresh cheese, butter and processed cheeses), especially when flavorings were present.

PROBIOTIC FORTIFICATION

Lactobacillus acidophilus or *bifidus* does not grow and survive in yogurt for a long period of time. A differential inoculation procedure has been developed where by *Lactobacillus acidophilus* is first inoculated into heat treated. Later, the regular yogurt cultures *Streptococcus thermophiles* and *Lactobacillus bulgaricus* are inoculated into the acidophilus growing yogurt mix. Dietetic fiber was introduced into the fruit base and then mixed with yogurt. Dietetic fiber enhances the population of *L. acidophilus*.

CASIENATE FORTIFICATION

Yoghurt fortification with caseinates, co-precipitate and blended dairy powders in low-fat yoghurt. These dairy products were characterised for pH, moisture, lactose, mineral and protein fractions. Yoghurts enriched with caseinates had higher viscosity and syneresis index than yoghurts based on concentrated skimmed milk fortified with co-precipitate or blended dairy products. One blended dairy product was tested to manufacture low-fat yoghurt on an industrial scale, yielding good rheological properties and lower cost than traditional enrichment with skimmed milk powder.

FIBER FORTIFICATION

Insoluble dietary fiber from five different sources (soy, rice, oat, corn and sugar beet) were used to fortify sweetened plain yogurt. In general, fiber addition led to lower overall flavor and texture scores. A grainy flavor and a gritty texture were intense in all fiber-fortified yogurts, except in those made with oat fiber. The evolution of organic acids during the fermentation and cold storage of control and oat-II-fiber-fortified yogurts showed a similar

pattern; only acetic and propionic acids were found in significantly higher amounts in the fiber- fortified product.

LIMITATIONS

WHO, FAO, Health Canada, and the Nestle Research Centre have acknowledged very few limitations regarding Food Fortification, which are investigated to rectify the limitations. Some are:

- Fortification of nutrients in foods may deliver toxic amounts of nutrients to an individual and also cause its associated side effects. Excessive fluoride intakes result in irreversible staining to the teeth.
- Fortification of foods with folic acid has been mandated in many countries solely to improve the folate status of pregnant women to prevent Neural Tube Defects. High blood concentrations of folic acid may decrease natural killer cell cytotoxicity, and high folate status may reduce the response to drugs used to treat malaria, rheumatoid arthritis, psoriasis, and cancer. Folate has a dual effect on cancer, protecting against cancer initiation but facilitating progression and growth of pre neoplastic cells and subclinical cancers.
- In many cases, the micronutrients added to foods in fortification are synthetic.
- In some cases, certain forms of micronutrients can be actively toxic in a sufficiently high dose, even if other forms are safe at the same or much higher doses. Retinol, the active form of Vitamin A, is toxic in a much lower dose than other forms, such as beta carotene.
- Menadione, a phased-out synthetic form of Vitamin K, is also known to be toxic.

Conclusion

Although it is recognized that food fortification alone will not combat this deficiency, it is a step towards reducing the prevalence of these deficiencies and their associated health conditions. Hence, research and technological application of food fortification will be a greater scope of compensation of deficient nutrients in food.

References

- [1] "Food Fortification Technology". Fao.Org. Retrieved 2016-03-30.
- [2] "Food Science | Educating Food Leaders for Over 100 years". Uoguelph. Ca. Retrieved 2016-03-30.
- [3] Fre De Ric Gaucheron, Iron Fortification In Dairy Industry, Trends In Food Science & Technology 11 (2000) 403–409

- [4] Helga Refsum; A. David Smith (August 2008). "Are We Ready For Mandatory Fortification With Vitamin B-12?" *The American Journal of Clinical Nutrition* **88** (2): 253–254. PMID 18689357
- [5] M.A. Drake, X.Q. Chen, S. Tamarapu and B. Leenanon, Soy Protein Fortification Affects Sensory, Chemical, And Microbiological Properties Of Dairy Yogurts, *Journal Of Food Science*, Volume 65, Issue 7, Pages 1244–1247, October 2000
- [6] Marta Guzmán-González, Federico Morais and Lourdes Amigo, Influence Of Skimmed Milk Concentrate Replacement By Dry Dairy Products In A Low-Fat Set-Type Yoghurt Model System. Use Of Caseinates, Co-Precipitate And Blended Dairy Powders, *Journal Of The Science Of Food And Agriculture*, Volume 80, Issue 4, Pages 433–438, March 2000
- [7] "Recommendations on Wheat and Maize Flour Fortification Meeting Report: Interim Consensus Statement" (PDF). *Who. Int.* Retrieved 2016-03-30.
- [8] S.H. Kang, J.U. Kim, J.Y. Imm, S. Oh, S.H. Kim, The Effects Of Dairy Processes And Storage On Insulin-Like Growth Factor-I (IGF-I) Content In Milk And In Model IGF-I-Fortified Dairy Products, *Doi:10.3168/Jds.S0022-0302(06)72104-X*
- [9] Wojciech Kolanowski, Jenny Weixbrodt, Sensory Quality of Dairy Products Fortified With Fish Oil, *International Dairy Journal* 17 (2007) 1248–1253.