

## BACTERIAL SPECIES ISOLATED FROM DIARRHOEIC CALVES AND ITS ANTIBIOTIC SENSITIVITY PATTERN

\*Manickam R<sup>1</sup> and Ponnusamy P<sup>2</sup>

<sup>1,2</sup>Department of Veterinary Microbiology, Veterinary College and Research Institute,  
Orathanadu, Thanjavur, Tamilnadu (India) – 614 625

E-mail: manickam75vet@gmail.com (\*Corresponding Author)

**Abstract:** This study was performed to identify various bacteria from feces of calves suffering from diarrhea, and to determine in vitro antimicrobial activity. Fecal samples were collected from 80 diarrheic calves and primarily tested for the presence of *Escherichia coli*, *Salmonella spp.* and *Staphylococcus aureus* using bacteriological examination, biochemical reactions. 60 bacterial isolates from the 80 fecal samples were identified using bacteriological and biochemical methods. *Escherichia coli* was considered to be the most frequent bacterium isolated numbering 22 (36.66%) followed by *Salmonella sp.* as the second most prevalent 11 (18.33%). Further isolates such as *Staphylococcus aureus* 8 (13.33%), *Enterococcus faecalis* 5 (8.33%) and *Shigella sp.* 3 (5.00%) were isolated and identified. Among the antibiotic sensitivity 55.0% were sensitive to Amikacin, 55% were sensitive to Ceftriaxone, 69.0% sensitive to Ciprofloxacin, 81.5% were sensitive to Kanamycin, and 75.5% were sensitive to Nalidixic acid (Table 4). Similarly majority of the bacterial isolates showed resistance to Ampicillin (75.0%), Amoxycillin (62.0%), Ceftriaxone (45.0%), Chloramphenicol (68.0%), Gentamicin (50.0%), Streptomycin (65.0%) and Tetracycline (74.0%).

**Keywords:** Bacteria, Calf diarrhea, *Escherichia coli*, *Salmonella* species - Antibiotic resistance.

### Introduction

Calf diarrhea caused by bacterial infection has a bad effect on the dairy industry all over the world when calves are reared intensively. It involves significant economic loss for labor and capital, calf mortality, loss in calf value and veterinary costs (Pereira et al., 2011; de Verdier et al., 2012). The pre-weaning mortality of dairy calves was estimated at 10.8% level with diarrhea responsible for more than half of those deaths (Pereira et al., 2011; Wu et al., 2010). Diarrhea caused by a variety of bacteria has been recognized as one of the most public clinical problems for calves worldwide. Among these bacteria *Eschirechia coli* (*E. coli*) as “white scour”, *Salmonella typhimurium* (*S. typhimurium*), *Clostridium perfringens* (*C. perfringens*) and *Staphylococcus aureus* (*St. aureus*) are believed to be the major microbial causes of diarrhea in calves (Hemashenpagam et al., 2009; Abdullah et al., 2013; Cho YI et al., 2010).

Antimicrobial agents are considered popular to fight diarrhea in calves. Nevertheless, their wide spectrum of activity, the emergence of microbial tolerance of different antimicrobial agents has become a well-known phenomenon, which represents a major concern (Hajipour et al., 2013). Resistance to antimicrobial agents was frequently occurred in *Salmonella* species and *E. coli* particularly in pre-weaned dairy calves (Izzo et al., 2011).

The frequent resistance of various microorganisms to the majority of antimicrobial agents is catching the attention of a great deal of awareness. The World Health Organization (WHO) commented on the severe threat formed by antibiotics-resistant bacteria in livestock and human health (Raffi et al., 2010). The frequent use of antibacterial agents has created the selective pressure to enhance the rising rates in antibiotic tolerance to different types of bacteria (El Zowalaty, 2012; Lee et al., 2013). Consequently, the microorganisms developed resistance against several types of antimicrobial agents to be one of the most important public health problems. In addition, the disadvantages of frequently used antimicrobial agents are not only the development of multiple drug resistance, but also adverse side effects (Huh and Kwon, 2011). As a result of antimicrobial resistance in the diarrhea of pre-weaned calves, there is an increasing attention in using alternative antimicrobial agent (Pereira et al., 2011; Hajipour et al., 2013). Consequently, there is an urgent need to find out an innovative approach and recognize new antimicrobial agents from natural and inorganic substances to develop the next generation of antimicrobial agents to control different microbial infections (Hajipour et al., 2013).

## **Materials and Methods**

### **Collection of samples:**

Rectal swabs were collected from (80) diarrheic calves using sterile cotton swabs and the samples were transferred directly to the laboratory in a separate clean sterile plastic bag, in an ice box and kept under complete aseptic condition without delay and subjected to bacterial culture, microscopic observation and antibiogram of *E.coli* and *Salmonella*.

### **Detection of bacterial fecal pathogens**

#### **Bacteriological examination:**

1. **Isolation and identification of *E. coli*:** All samples were inoculated into tubes of freshly prepared nutrient broth and incubated aerobically at 37°C over night, followed by subculturing onto MacConkey agar and Eosin methylene blue agar plates for 24-48 hours at 37°C. Lactose positive colonies were confirmed as *E. coli* according to Gershwin (1990); Koneman *et al.*, (1992) and Quinn *et al.*, (1994). Suspected colonies grown were picked on

nutrient agar slopes and incubated at 37°C for 24 hours, then kept in refrigerator at 4°C for further identifications according to Edwards and Ewing (1972).

## **2. Isolation and identification of Salmonella**

All samples were inoculated into tubes of Selenite-F and Tetrathionate broths and streaked out onto MacConkey and brilliant green agar after overnight incubation at 37°C. Suspected colonies were subjected to biochemical testing according to Adyin *et al.*, (2001) and Echeita *et al.*, (2002). Slide agglutination test was used for identification according to the Kauffmann-White Schema (Zahraei *et al.*, 2007; Nori and Thong, 2010). Finally identification of *Salmonella typhimurium* was done according to Waltner-Toews *et al.*, (1986).

## **3. Isolation and identification of Staphylococcus**

Isolation of presumptive *S. aureus* from the samples was performed using Mannitol salt agar (MSA). The plates were incubated aerobically at 37°C for 18 h – 24 h. Consequently, the characteristic *S. aureus* colonies that were yellow in color from MSA plate were further purified by sub-culturing onto MSA plates and the plates were incubated aerobically at 37°C for 18 h–24 h. These isolates were retained for further bacterial identification.

### **Microscopic Study by Staining Method**

The microorganisms were isolated from suspected cases of fecal samples, and then stained with Gram's staining techniques (Merchant and Packer, 1967).

### **Identification of bacterial isolates by using specific biochemical tests**

After culture characterization of *E. coli*, *Salmonella* spp. and *St. aureus* on different culture media as MacConkey agar, EMB agar, Blood agar, SS agar and Mannitol salt agar, all the isolates were identified by various biochemical tests including Catalase, Oxidase, Triple sugar iron (TSI) agar slant reaction, Voges–Proskauer (VP), Methyl-Red (MR), Indole, Motility indole and urease (MIU) test (Merchant and Packer, 1967).

### **Antibacterial Sensitivity Pattern of the Isolated Salmonella and E. coli**

The overnight nutrient broth cultured *Salmonella* isolates were poured on SS agar and spread uniformly with the help of sterile glass spreader. Antibacterial discs were applied aseptically to the surface of the plate at an appropriate distance with the help of sterile forceps and incubated at 37°C for 24 hours, aerobically. Antibiotic sensitivity pattern of isolated *E. coli* and *Salmonella* were performed against 11 commonly used antibiotics belonging to different groups (Bauer *et al.*, 1966) procured from HiMedia Pvt. Ltd. The antimicrobial agents tested were the following; Amikacin (AK) , Ampicillin (AMP), Ceftriaxone (CTR), Chloramphenicol, Ciprofloxacin (CIP), Gentamicin (GEN), Kanamycin, Nalidixic acid (NA),

Streptomycin (S) and Tetracycline (TE). Sensitivity of the isolated bacterial strains to selected antibiotics and sulfonamides was tested using the Kirby-Bauer disk diffusion method on Mueller-Hinton medium in accordance with accepted international norms (CLSI, 2011). The results were read and interpreted based on the diameter of the zone of inhibition, with the strains designated as resistant (R), of intermediate sensitivity (I) or sensitive (S).

## Results

### Characterization of bacterial fecal pathogens

Isolation and identification of *E. coli*, *Salmonella* spp., and *St.aureus* were carried out by using morphological, biochemical and genotypic characteristics. A total of 60 bacterial isolates were obtained from the 80 fecal samples. *E. coli* was considered the most frequent bacterium isolated (36.66%) followed by *Salmonella* sp. as the second most prevalent (18.33%) (Table 3). Further isolates such as *Klebsiella* spp. (18.33%), *St. aureus* (13.33%), *Enterococcus faecalis* (8.33%) and *Shigella* sp. (5.00%) were isolated respectively. In the current study, the sample found positive for *E. coli* gives a positive reaction to lactose fermentation on MacConkey agar plate, metallic green sheen colonies on EMB plates and yellowish green colonies on Brilliant Green Agar (BGA). The sample gives a positive reaction for *Salmonella* producing negative reaction to lactose fermentation on MacConkey agar plate, Opaque, transparent and pale colonies with black center were produced on SS agar and pink color colonies on Brilliant green agar. Additionally, the sample gives a positive reaction for *St. aureus* producing yellowish colonies on tryptose soya agar and hemolysis on blood agar and yellow colonies on Mannitol salt agar (Table 1).

The various isolates of *E. coli*, *Salmonella* spp. and *St. aureus* demonstrated identical results in various biochemical assays including sugar fermentation, Triple Sugar Iron (TSI) slant, Motility Indole Urease (MIU) test, Indole, Methyl Red and Voges-Proskauer Test (MR-VP), citrate utilization tests and Coagulase test. *St. aureus* produce acid but no gas by fermenting various sugars and gave positive reaction to coagulase, catalase and methyl red tests but negative reaction to Indole and Voges Proskeur test (Table 2).

**Table 1:** Identification of isolated microbial pathogens by cultural properties

Culture media utilized	Detection (Colony morphology)		
	<i>E.coli</i>	<i>Salmonella spp.</i>	<i>St.aureus</i>
Nutrient agar	Smooth, rounded, white to grayish colony with unusual putrid odor	Small, circular and smooth colonies	Growth of circular, small, smooth, convex, and golden yellowish colonies

Blood agar	Demonstrate haemolysis	Demonstrate haemolysis	Demonstrate haemolysis
Mac Conkey agar	Rosy pink lactose fermenter colonies	Colorless, pale, translucent colony	No growth (-)
Salmonella, Shigella (SS) agar	Pink color colony	Translucent colorless smooth colony with black center	No growth (-)
Mannitol salt agar	No growth (-)	No growth (-)	Yellow colonies
Eosin-Methylene Blue (EMB) agar	Moist circular colonies with dark centers, yellow green metallic sheen	Light purple to colorless	No growth (-)

**Table 2:** Biochemical characteristics of *E. coli* and *Salmonella* from diarrheic calves

isolated bacteria	Indole production test	Methyl-red test	Voges-Poskauer reaction	Citrate utilization test	MIU test	TSI Test	Hydrogen sulphide
<i>E. coli</i>	+	+	-	-	+	Butt-Y Slant-Y	-
<i>Salmonella spp</i>	-	+	-	-	-	Butt-Y Slant-R	+
<i>Klebsiella spp</i>	-	-	+	+	-	Butt-Y Slant-Y	-
<i>St.aureus</i>	-	+	+	+	-	Butt-R Slant-R	-
<i>Enterococcus faecalis</i>	-	-	+	-	-	Butt-R Slant-R	-
<i>Shigella spp.</i>	-	+	-	-	-	Butt-Y Slant-R	-

**Table 3:** Frequency of bacterial species isolated from calves suffering from diarrhea

Bacterial isolates	Number of isolates	Percentage (%) of isolates
<i>Escherichia coli</i>	22	36.66
<i>Salmonella spp</i>	11	18.33
<i>Klebsiella spp</i>	11	18.33
<i>St. aureus</i>	8	13.33
<i>Enterococcus faecalis</i>	5	8.33

<i>Shigella spp</i>	3	5.00
Total	60	99.98

### Antibiotic sensitivity test of various isolates of *E. coli* and *Salmonella*

Among the antibiotic sensitivity 55.0% were sensitive to Amikacin, 55% were sensitive to Ceftriaxone, 69.0% sensitive to Ciprofloxacin, 81.5% were sensitive to Kanamycin, and 75.5% were sensitive to Nalidixic acid (Table 4). Similarly majority of the bacterial isolates showed resistance to Ampicillin (75.0%), Amoxicillin (62.0%), Ceftriaxone (45.0%), Chloramphenicol (68.0%), Gentamicin (50.0%), Streptomycin (65.0%) and Tetracycline (74.0%).

**Table 4:** Antibiotic sensitivity of bacterial species isolated from calves suffering from diarrhea

Antibiotics	Sensitive (%)	Intermediate (%)	Resistant (%)
Amikacin	55.00	2.50	42.50
Ampicillin	25.00	--	75.00
Amoxicillin	35.00	3.00	62.00
Ceftriaxone	55.00	--	45.00
Chloramphenicol	30.50	1.50	68.00
Ciprofloxacin	69.0	--	31.00
Gentamicin	47.75	2.25	50.00
Kanamycin	81.50	--	18.50
Nalidixic acid	75.50	--	24.50
Streptomycin	35.00	--	65.00
Tetracycline	26.00	--	74.00

## Discussion

### Identification of bacterial fecal pathogens

In the current investigation, various types of bacteria (*E. coli*, *Salmonella spp.*, *Klebsiella spp.*, *St. aureus*, *Enterococcus faecalis* and *Shigella spp.*) were isolated from a total of 80 fecal specimens collected from from pre-weaned calves suffering from diarrhea. 22 samples were identified as *E. coli* which appeared as rosy pink lactose fermenter colonies on MCA plate, moist circular colonies with dark yellow centers, green metallic sheen on EMB plates and yellowish green colonies on the BGA. Eleven samples were identified phenotypically as *Salmonella spp.* because the organism on MCA plate gave colorless, pale, translucent colony with negative reaction to lactose fermentation, translucent colorless smooth colonies with black center on S-S agar, pink color colonies on BGA and pink colour colony with black center on DCA. Eight samples were found positive for *St. aureus* producing yellow pigmented golden colonies on Trypticsoy agar, yellow colour colony on Mannitol salt agar

and hemolysis on Blood agar. Dissimilarities in the colony shape appeared by the isolates may be as a result of losing or obtaining some properties by the transfer of host or choice of host tissue as noticed by Abdullah et al., (2013).

Various biochemical tests, including Indole, Voges-Poskauer, Methyl-red, Citrate, TSI, MIU and Coagulase were used to distinguish between different isolates of bacteria in this study. *E. coli* gave positive reactions to Indole and Methyl-red tests, but negative to Voges- Poskauer, Citrate and Coagulase tests. *St. aureus* gave positive reactions to Coagulase and Methyl red tests, however negative reaction to Indole, Voges Proskeur, Citrate and MIU tests was reported. Moreover, *Salmonella spp.* gave positive results to Methyl-red and MIU tests while negative to the other tests. To date, the real reasons for which the appearance of an identical results in biochemical reactions of the three groups of recognized isolates were not obvious. There is no doubt that nearly all bacterial isolates in the current investigation have several common genetic materials which might be responsible for the appearance of similar type of biochemical reaction as reported by Pandey et al., (1979) and Honda et al., (1982).

#### **Antibiotic resistance and antimicrobial effect of various antibiotics**

The present study is to evaluate the effect of various antibiotics on *E. coli*, *Salmonella* and *St. aureus* isolated from feces of calves suffering from diarrhea. Therefore, the antimicrobial sensitivity test of three various types of bacterial isolates to 11 different antimicrobial agents were studied. The sensitivity investigation demonstrated that the majority of the *E. coli*, *Salmonella spp.* and *St.aureus* were tolerant to ampicillin, tetracycline, amoxicillin and chloramphenicol. Similar results were obtained by Abdullah et al., 2013; Ahmad et al., 1986; Edrington et al., 2004; Nazir, (2007) who stated that calf isolates were resistant to ampicillin, amoxicillin, erythromycin and gentamicin (Nazir, 2007). These findings were somewhat different with the results of Guerra et al., 2006 and Joon and Kaura, (1993) who stated that most of the microorganisms isolated from calves suffering from diarrhea were highly susceptible to chloramphenicol and tetracycline and moderately sensitive to ampicillin and amoxicillin (Joon and Kaura, 1993). The variation in the susceptibility of antimicrobial agents against the fecal isolates may be due to the outcome of selection and also the random apply of antibiotic in various disease stages to different animal species. The appearance of multidrug-resistant bacteria is documented as an important problem for public health worldwide. Consequently, treatment is expensive and needs prolonged time.

## References

- [1] Abdullah M, Akter MR, Lutful Kabir SM, Abu Sayed Khan, M, Abdul Aziz MS (2013) Characterization of Bacterial Pathogens Isolated from Calf Diarrhea in Panchagarh District of Bangladesh. *J Agric Food Tech* 3:8-13.
- [2] Adyin, F.; Mmur, S.; Gokce, G.I.; Genc, O. and Guler, M.A. (2001): The isolation and identification of bacteria and parasites from diarrhea calves in Kars District. *Kafkas universitesi veteriner Fakultesi, Dergisi*. 7, (1): 7.
- [3] Ahmad R, Amin M, Kazmi SE (1986) Studied on the bacterial causes of calf mortality. *Pak Vet J* 6: 116-118.
- [4] Bauer, A.W., Kirby, W.M.M., Sherris, J.C. and Turck, M. (1966). Antibiotic susceptibility testing by a standardized single disk method. *Amer. J. Clin. Pathol.*, 45: 493-496.
- [5] Cho YI, Kim WI, Liu S, Kinyon JM, Yoon KJ (2010) Development of a panel of multiplex real-time polymerase chain reaction assays for simultaneous detection of major agents causing calf diarrhea in feces. *J Vet Diagn Invest* 22: 509-517.
- [6] CLSI (Clinical and Laboratory Standards Institute, 2011) Performance Standards for Antimicrobial Susceptibility Testing; Twenty-First Informational Supplement 31: 68-78.
- [7] de Verdier K, Nyman A, Greko C, Bengtsson B (2012) Antimicrobial resistance and virulence factors in *Escherichia coli* from Swedish dairy calves. *Acta Vet Scand* 54: 2.
- [8] Echeita, MA.; Herrera, S.; Garaizar, J. and Usera, MA. (2002): Multiplex PCR based detection and identification of the most common *Salmonella* second-phase flagellar antigen. *Res. Microbiol.* 153: 107-113.
- [9] Edrington TS, Hume ME, Loofer ML, Schultz CL, Fitzgerald AC, (2004) Variation in the faecal shedding of *Salmonella* and *E. coli* O157:H7 in lactating dairy cattle and examination of *Salmonella* genotypes using pulsed-field gel electrophoresis. *Lett Appl Microbiol* 38:366-372.
- [10] Edwards, P.R. and Ewing, W.H. (1972): Identification of Enterobacteriaceae. Burgess publ Co. Minnece polis, Minnesota, p.103-104.
- [11] El Zowalaty ME (2012) Alarming trend of antibiotic resistance in *Pseudomonas aeruginosa* isolates. *Journal of Pure and Applied Microbiology* 6: 175–183.
- [12] Gershwin, L.J. (1990): The physiochemical and biological basis of immunity. In: Biberstein, E.L., Zee, Y.C. (Eds.): Review of Veterinary Microbiology. Blackwell Scientific Publications, Boston, USA. 29–30.



- [13] Guerra B, Junker E, Schroeter A, Helmuth R, Guth BE, (2006) Phenotypic and genotypic characterization of antimicrobial resistance in *Escherichia coli* O111 isolates. *J Antimicrob Chemother* 57: 1210-1214.
- [14] Hajipour MJ, Fromm KM, Ashkarran AA, de Aberasturi DJ, de Larramendi IR (2013) Antibacterial properties of Nanoparticles. *Trends in Biotechnology* 31: 61-62.
- [15] Hemashenpagam N, Kiruthiga B, Selvaraj T, Panneerselvam A (2009) Isolation, Identification and Characterization of Bacterial pathogens causing Calf Diarrhea with special reference to *Escherichia coli*. *The Internet Journal of Microbiology* 7(2).
- [16] Honda T, Arita M, Takeda Y, Miwatani T (1982) Further evaluation of the Biken test (modified Elek test) for detection of enterotoxigenic *Escherichia coli* producing heat-labile enterotoxin and application of the test to sampling of heat-stable enterotoxin. *J Clin Microbiol* 16: 60-62.
- [17] Huh AJ, Kwon YJ (2011) "Nanoantibiotics": a new paradigm for treating infectious diseases using nanomaterials in the antibiotics resistant era. *J Control Release* 156: 128-145.
- [18] Izzo M, Mohler V, House J (2011) Antimicrobial susceptibility of *Salmonella* isolates recovered from calves with diarrhoea in Australia. *Aust Vet J* 89: 402-408.
- [19] Joon DS, Kaura YK (1993) Isolation and characterization of Enterobacteria from diarrhoeic and non-diarrhoeic calves. *Indian J Anim Sci* 63: 373-383.
- [20] Koneman, E.W.; Allen, S.D.; Dowell, V.R.; Janda, W.H. and Sommers, H.M. (1992): *Color atlas and Textbook of Diagnostic Microbiology*. 4th Ed., J.B.Lippincott CO., New York.
- [21] Lee CR, Cho IH, Jeong BC, Lee SH (2013) Strategies to minimize antibiotic resistance. *Int J Environ Res Public Health* 10: 4274-4305.
- [22] Merchant IA, Packer RA (1967) *Veterinary Bacteriology and Virology*. (7th Edn) Ames, Iowa, Iowa State University Press 752p.
- [23] Nazir KH (2007) Plasmid profiles and antibiogram pattern of *Escherichia coli* isolates of calves feces and diarrhegenic stool of infants. *Journal of Bangladesh Society of Agricultural Science and Technology* 4(1&2):149-152.
- [24] Nori, EM. And Thong, KL. (2010): Differentiation of *Salmonella enterica* based on PCR detection of selected somatic and flagellar antigen. *Afr. J. Microbiol. Res.* 4(9): 871-879.
- [25] Pandey PN, Thaphyal DC, Sharma SN (1979) Enterotoxigenicity of some *Escherichia coli* isolates. *Indian J Anim Research* 13: 1-4.

- [26] Pereira RV, Santos TM, Bicalho ML, Caixeta LS, Machado VS (2011) Antimicrobial resistance and prevalence of virulence factor genes in fecal *Escherichia coli* of Holstein calves fed milk with and without antimicrobials. *J Dairy Sci* 94: 4556-4565.
- [27] Quinn, P.J.; Carter, M.E.; Markey, B.K. and Carter, G.R. (1994): *Clinical Veterinary Microbiology*. Mosby. Yearbook Europe Limited.
- [28] Raffi M, Mehrwan S, Bhatti TM, Akhter JI, Hameed A, (2010) Investigations into the antibacterial behavior of copper nanoparticles against *Escherichia coli*. *Annals of Microbiology* 60: 75-80.
- [29] Waltner-Toews, D.; Martin, S.W. and Meek, A.H. (1986): An epidemiological study of selected calf pathogens on Holstein dairy farms in southwestern Ontario. *Canadian Journal of Veterinary Research*, 50: 307–313.
- [30] Wu G, Mafura M, Carter B, Lynch K, Anjum MF (2010) Genes associated with *Escherichia coli* isolates from calves with diarrhea and/or septicaemia. *Vet Rec* 166: 691-692.
- [31] Zahraei, T.; Tadjbakhsh, H.; Atashparvar, N.; Nadalian, MG. and Mahzounieh, M.R. (2007): Detection and identification of *Salmonella* Typhimurium in bovine diarrhoeic fecal samples by immunomagnetic separation and multiplex PCR assay *Zoonosis Public Health*, 54: 231-236.