

**IN VITRO ANTIFUNGAL ACTIVITY OF ESSENTIAL OILS OF
AROMATIC PLANTS OF THE SENEGALESE FLORA ON
COLLETOTRICHUM GLOEOSPORIOIDES (PENZ.) PENZ AND SACC.,
THE CAUSAL AGENT OF MANGO ANTHRACNOSE**

Maodo Malick CISSE¹, Nalla MBAYE¹ and Papa M. Diédhiou^{2*}

¹Département de Biologie Végétale, Université Cheikh Anta DIOP, Dakar BP 5005, Sénégal

²UFR des Sciences Agronomiques d'Aquaculture et des Technologies Alimentaires
(UFR S2ATA), Université Gaston Berger de Saint-Louis (UGB), Saint-Louis, Sénégal

E-mail: anifane@gmail.com, papa-madiallacke.diedhiou@ugb.edu.sn

(*Corresponding author)

Abstract: Mango production has gained considerable economic importance in Senegal. However, it suffers from major phytosanitary constraints the main of which is anthracnose caused by *Colletotrichum gloeosporioides*. The control of anthracnose relies mainly on synthetic fungicides with the known negative impact on the environment as well as on consumer health. Alternatives to synthetic fungicides like essential oils were reported to have great potential on fungal pathogens. In this study, the efficacy of essential oils of plants such as those from *Mentha piperita*, *Ocimum basilicum*, *Melaleuca quinquenervia* and *Eucalyptus camaldulensis* was studied *in vitro* on mycelial growth, spore germination and spore production of *Colletotrichum gloeosporioides*. The essential oils of *M. piperita* and *Ocimum basilicum* exhibited the best results. *M. piperita*' essential oil inhibited mycelial growth and spore production by more than 50% at 1000 ppm and suppressed them at 3000ppm. The essential oil of *M. piperita* was able to suppress spore germination at 500 ppm while for that of *O. basilicum* a similar result was observed at 1500 ppm. These extracts could offer potential for use in biological control.

Keywords: *Colletotrichum gloeosporioides*, Essential oils, Biological control, Senegal.

Introduction

In Senegal, the mango sector employs nearly 30,000 people and generates a turnover of more than 10 billion FCFA per year (Asepex, 2012). Various constraints such as the high pressure of pathogens including *C. gloeosporioides*, responsible for anthracnose, must however be lifted to increase its profitability. This disease is one of the major post-harvest constraints and causes also huge damages during production of mango in the world (Pardo De la Hoz *et al.*, 2016). In Senegal, the losses can reach 42% in the Niayes zone (Mbaye, 2006) and go up to 90% in Casamance during the rainy season (Diédhiou *et al.*, 2014). Disease control rely to date, on the use synthetic fungicides. However, a frequent use of pesticides often increases the risk of

Received April 2, 2020 * Published June 2, 2020 * www.ijset.net

residues in mangoes, environmental pollution and threat to human and animal health. In response to the growing consumer anxiety and increased demand in the global marketplace for safer food, research is developing alternative control methods such as the use of essential oils. Many studies on controlling fungal diseases with essential oils and plant extracts such as those of Padman and Janardhana (2012), Pandey *et al.* (2012); Gautam *et al.* (2013), Ba (2014) have already been published. The essential oils of *M. piperita*, *Ocimum basilicum*, *Melaleuca quinquenervia* were suspected to have antifungal properties because of their chemical composition Carretto *et al.* (2010), Edris and Farrag (2003), Funery *et al.* (2006).

In this study, the efficacy of the essential oils of *M. piperita*, *O. basilicum*, *M. quinquenervia* and *Eucalyptus camaldulensis* was tested *in vitro* on *C. gloeosporioides*.

Material and methods

Fungal material

Strains of *Colletotrichum gloeosporioides* were isolated from infested mangoes of the variety Kent from Casamance.

Essential oils

The steam extraction method was used to obtain the essential oils from the biomass of the several plant species targeted. Therefore, the whole plant of *Mentha piperita*, and the leaves and flowers of *O. basilicum*, were submitted to extraction after 10 days of drying at room temperature. As for *Eucalyptus camaldulensis* and *Melaleuca quinquenervia*, their essential oils were extracted from fresh leaves.

After extraction, the essential oils were stored in tinted vials at room temperature in the laboratory for protection from light.

***In vitro* activity of essential oils on mycelial growth**

Preparation of fungal cultures

The essential oils were incorporated into the PDA (Potato Dextrose Agar) culture medium after autoclaving at a temperature of 121 °C, at a pressure of 1 bar for 25 minutes and then poured into Petri dishes of 9 cm in diameter at a rate of 20 ml. per box. For each concentration of essential oils, 3 PDA plates were inoculated with 3 mm diameter of agar disc colonized by a pure culture of *Colletotrichum gloeosporioides* aged of 7 days. An incubation period of 8 days was considered because it corresponded to time interval taken by the fungus to fill the non-treated petri dishes.

Preparation of different concentrations

The essential oils are first mixed with Tween 80 (0.1%) before being incorporated in the PDA culture medium after sterilization and stirred for a few minutes to allow good homogenization. Concentrations ranging from 1000 to 10000 ppm were tested for all the essential oils tested.

Evaluation of mycelial growth

Mycelial growth was assessed every 48h by calculating the average colony diameter of perpendicular measurements passing through the middle of the fungal colony. Three repetitions are performed for each concentration. Growth inhibition rates (GIR) were determined according to the formula from Doumbouya *et al.* (2012):

$GIR (\%) = (T-E) * 100 / T$; where

GIR = Growth inhibition rate

T = Average diameter of fungal colony in the control treatment (in cm)

E = Average diameter of fungal colony in treated plates (in cm)

The LD50s were subsequently deduced on the basis of the Probit method of Finney (1941) by using the `dose.p` function of the R software MASS package.

Activity of essential oils on the production of spores of *C. gloeosporioides*

The fungal strain growing in the Petri dishes containing the media amended with the essential oils, 10 mL of sterile distilled water with Tween 80 (0.1%) were added. Three minutes later, the mycelium is scrapped using a scalpel blade and then filtered twice to have a spore suspension free from mycelial fragments. The resulting spore suspension was diluted 100-fold with sterile distilled water. Using a micropipette, 50 μ L of this suspension were taken and deposited on a hemocytometer. The number of spores was counted under an optical microscope at X 400 magnification. Five repetitions were made for each concentration as well as with the control treatment. Inhibition of spore production was determined by the formula proposed by Doumbouya *et al.* (2012):

$ISP = (SCo0-SCoT) * 100 / SCo0$

ISP = inhibition of sporulation performance (in %)

SCo0 = spore concentration in control treatment

SCoT = spores concentration in the treatment considered

The LD50s are subsequently deduced using the Finney Probit method (1941) using the `dose.p` function of the MASS package of the R software.

Activity of essential oils on spore germination of *C. gloeosporioides*

In a seven-day old culture dish of *C. gloeosporioides*, 10 ml of sterile distilled water with Tween 80 solution (0.1%) was added. The mycelium was collected from petri dish and put into a beaker containing 100 ml sterile distilled water. The whole was shaken during 3 min before filtration over a 4 layers gaze clothe to obtain a spore suspension. The spore suspension was diluted 100 times with sterile distilled water and 50 µl of the suspension were mixed with 50 µl of essential oil previously mixed with 0.1% Tween. A drop of the mixture was then deposited between a slide and coverslip. The preparations were placed in Petri dishes containing moistened tissue to maintain a high relative humidity.

The observations were made every 12 hours for 2 days and three repetitions are performed for each concentration of essential oil which ranged from 100 to 1500 ppm. The number of germinating spores was counted as based on the initiation of the germ tube. The percentage of germination in the different treatment was confronted to spore germination in the control to assess inhibition performance of the essential oils using the formula proposed by Doumbouya *et al.*, (2012):

$$GR = \text{NGSpT} / \text{NGSpC} * 100$$

GR = germination rate

NGSpT = Number of germinating spores in considered treatment

NGSpC = Number of germinating spores in control treatment

The LD50s are subsequently deduced on the basis of the Probit method of Finney (1941) using the dose.p function of the R software MASS package.

Statistical analysis

The data was processed with R.3.2.3 software (R Core Team, 2015). A two-factor analysis of variance (ANOVA) was performed using the aov function of the agricolae package (de Mendiburu, 2015).

Results

*Activity of essential oils on mycelial growth of *C. gloeosporioides**

Mentha piperita had a high efficacy at inhibiting mycelial growth of the fungus with 77% at 1000 ppm and total inhibition at 3000ppm. It is followed by *Ocimum basilicum* where total inhibition was obtained at 6000ppm (Figure 1). The essentials oils of Melaleuca and Eucalyptus were the less effective but still achieve at least 80% inhibition of mycelial growth at 10 000 ppm.




Figure 1: Activity of essential oils on the mycelial growth of *Colletotrichum gloeosporioides* (n = 120, P = 2e-16).

EC = Eucalyptus camaldulensis, MFF = Melaleuca quinquenervia (fresh leaves), MFS = Melaleuca quinquenervia (dry leaves), MP = Mentha piperita, OB = Ocimum basilicum

Activity of essential oils on the sporulation of *C. gloeosporioides*

The essential oils inhibited significantly the sporulation of the pathogen. On that parameter also, the mint oil was still the most effective with a reduction of 60% of fungal sporulation at 1000 ppm and total inhibition achieved at 3000ppm (Figure 2). With the basilic oil, fungal sporulation was reduced at 40% at 1000 ppm and with 6000ppm, sporulation was inhibited completely. The essentials oils of Melaleuca and Eucalyptus were the less effective but still achieve total inhibition of fungal sporulation at 10 000 ppm.




Figure 2: Activity of essential oils on the production of spores of *C. gloeosporioides* (n = 120, P = 2e-16).

EC = Eucalyptus camaldulensis, MFF = Melaleuca quinquenervia (fresh leaves), MFS = Melaleuca quinquenervia (dry leaves), MP = Mentha piperita, OB = Ocimum basilicum

Activity of essential oils on spore germination of *C. gloeosporioides*

Spore germination of *C. gloeosporioides* was about 70 to 80% without treatment (Figure 3). It was reduced down to 20% and 30% through treatment with respectively 100 ppm of essential oil of *Mentha piperita* and *Ocimum basilicum*. Spore germination was further totally inhibited at 500ppm of the essential oil of *M. piperita* while with *O. basilicum* total suppression of spore germination was observed at 1500ppm. The essential oil of *E. camaldulensis* reduced spore germination down to $21.66 \pm 14.37\%$ at 1500 ppm.




Figure 3: Activity of essential oils on the germination of spores of *Colletotrichum gloeosporioides* (n = 75, P = 2e-16)

EC = *Eucalyptus camaldulensis*, MFF = *Melaleuca quinquenervia* (fresh leaves), MFS = *Melaleuca quinquenervia* (dry leaves), MP = *Mentha piperita*, OB = *Ocimum basilicum*

Determination of the lethal concentration of essential oils

For the essential oil of *Mentha piperita* the LC50 for spore germination was 48.12 ppm (Table 1). The value raised up to 156.34 ppm for mycelial growth and was about 749.12 ppm for spore production. For *O. basilicum*, the LC50 reached 85.02 ppm for spore germination and was at 156.34 for mycelial growth, while 2242.93 ppm was the LC50 for inhibition of sporulation.

Table 1: Lethal Concentration 50 (LC50) in ppm of Essential Oils on Measured Parameters

| | Spore germination | Mycelial growth | Sporulation |
|---------------------------------|-------------------|-----------------|-------------|
| <i>Eucalyptus camaldulensis</i> | 376,12 | 3752,42 | 5762,79 |
| <i>Melaleuca quinquenervia</i> | 322,45 | 416,57 | 4067,06 |

| | | | |
|--------------------------------|--------|---------|---------|
| <i>Melaleuca quinquenervia</i> | 594,84 | 1780,81 | 4899,18 |
| <i>Mentha piperita</i> | 48,12 | 156,34 | 749,12 |
| <i>Ocimum basilicum</i> | 85,02 | 2243,47 | 2142,93 |

Discussion

Essential oils could be an important source of solutions towards the environment while being consumer and user friendly alternatives for chemical fungicides. Their performance against the mycelial growth, spore germination and spore production of *C. gloeosporioides*, was very important. Among the 5 tested oils, the antifungal effect of *M. piperita* essential oil was the effective. It allowed to obtain a total inhibition of mycelial growth and spore production at 3000 ppm while spore germination was totally suppressed at 500 ppm. The essential oil of *M. piperita* is composed mostly of menthol which is a monoterpene (Mahboubi and Kazempour, 2014), menthone and limonene (Iskan *et al.*, 2002). These molecules are suspected of causing disruption of cell membranes of fungi (Ghosh *et al.*, (2005), which could be one of the mechanisms of action leading to the observed performance of the essential oil of *M. piperita*. The menthol and menthone are also speculated to initiate an alteration of cell metabolism of fungi that leads to complete destruction (Zani *et al.*, 1991). Those 2 mechanisms linked to the active molecules in the essential oil of *M. piperita* could be at the origin of the efficiency observed in mycelial growth as well as sporulation and spore germination of *C. gloeosporioides*.

The essential oil of *O. basilicum* was highly effective against the pathogen with 100% inhibition of mycelial growth and spore production at 6000 ppm and spore germination at 1500ppm. A strong inhibitory activity of this essential oil of *O. basilicum* on mycelial growth, spore germination and spore production of plant pathogens such as *Phytophthora capsici*, *Phytophthora dreschleri* and *Phytophthora melonis* has already been reported (Amini, *et al.*, (2016). The essential oil of *O. basilicum* was reported to contain of more than 20 chemical compounds including terpenic compounds such as linalool (Amini *et al.*, 2016), 1,8 cineol and methyl eugenol (Govindarajan *et al.*, 2013). It has antimicrobial properties (Sokovic *et al.*, 2010, Rao *et al.*, 2011, Matiz *et al.*, 2012), antioxidant activity (Trevisan *et al.*, 2006) and antifungal futures (Edris and Farrag, 2003, Amini *et al.*, 2016). Some terpenic compounds are known to be able to penetrate into the phospholipid double layer of the membrane of the cell

and induce its rupture. The cytoplasmic contents are thus discharged outside the cell involving its destruction (Tsuchiya *et al.*, 1996). They are also reported to induce a chemo-osmotic disruption and intra-cytoplasmic potassium leakage, followed by release of nucleic acids, ATP, and inorganic phosphate (Mokaddem, 2011). Those mechanisms might be at the basis of the efficiency of the essential oil of *O. basilicum* on the inhibition of fungal growth but also on spore germination and the sporulation of fungi.

The essential oil of *M. quinquenervia* is composed of various chemical compounds including terpinene-4-ol and 1,8-cineol, which have a broad spectrum of antimicrobial activity, including antibacterial, antiviral and antifungal properties (Funery *et al.*, 2006, Shelton *et al.*, 2004). According to Doumbouya *et al.* (2012), the essential oil of *M. quinquenervia*, causes total inhibition of mycelial growth, spore germination and spore production of *C. gloeosporioides* at 8500 ppm. The antifungal capacity of this essential oil on *C. gloeosporioides* was confirmed by the present study showing but total inhibition of the fungus was obtained at a higher concentration. Variability in the composition and concentration of the essential oils of the same species according to location and time of harvest of biomass as well as the genotype of fungi tested might be the cause of this variation.

With the essential oil of *Eucalyptus camaldulensis* spore germination of *C. gloeosporioides* was reduced down from only 80% to 21.66 % at a concentration of 1500 ppm, which ranked medium among the tested oils. Its activity on mycelial growth and spore production where the respective the least effective. *Eucalyptus camaldulensis* is one of the most widespread plant species worldwide (Gil *et al.*, 2010). Its oil contains 1,8 cineol, p-cimene (Shahwar *et al.*., 2012), ethanol, eucalyptol and carvacrol (Akin *et al.*, 2012). The essential oil of *E. camaldulensis* had a moderate efficiency against *C. gloeosporioides* despite the insecticidal, nematocidal (Shahwar *et al.*, 2012), antibacterial and fungicidal properties of its chemical compounds (Bamayi *et al.*, 2004).

The essential oils of *M. piperita* and *Ocimum basilicum* are rich in terpene molecules such as 1,8-cineol and terpinene-4-ol, (Funery *et al.*, 2006) (Akin *et al.*, 2012). Arras and Usai (2001) showed that these aromatic molecules increase the cell permeability, induce a degradation of nucleic acids and cause an inhibition of mitochondrial energy metabolism (Lambert *et al.*, 2001) in microorganisms. These alterations result in major disruption of physiological and biochemical processes in the cell leading therefore to the inhibition of the microorganisms (Yoshimura *et al.*, 2010).

References

- [1] **Akin M. and Aktumsek A. 2012.** Antibacterial activity and composition of the essential oils of *Eucalyptus camaldulensis* Dehn. and *Myrtus communis* L. growing in Northern Cyprus. *Afr. J. Biotechnol.* 9.
- [2] **Amini J, Fahrang V, Taimoor J, Nazemi J. 2016.** Antifungal Effect of Plant Essential Oils on Controlling *Phytophthora* Species. *The Plant Pathology Journal*, 32 (1):16-24.
- [3] **Arras G, Usai M. 2001.** Fungitoxic activity of 12 essential oils against four postharvest citrus pathogens: chemical analysis of thymus capitatus oil and its effect in subatmospheric pressure conditions. *J Food Prot.*; 64(7): 1025-9.
- [4] **Asepex, 2012.** Can smallholder producers ever succeed in export supply chains? Case study: Senegal Mango sector. Ministère du Commerce, de l'Industrie et de l'Artisanat. 18P
- [5] **Ba A. 2014.** Utilisation d'extraits de plantes médicinales pour le contrôle in vitro de *Colletotrichum gloeosporioides* Penz., agent responsable de l'antracnose de la mangue. Mémoire de Master, Biologie Végétale, Fac. Sciences, Université Cheikh Anta Diop de Dakar 55 P.
- [6] **Bamayi H, Kolo L, Okogun II, Ijah UJJ. 2004.** The Antimicrobial activities of methanol extracts of *Eucalyptus camaldulensis* and *Terminalia catappa* against some pathogenic microorganisms. *Journal of Nigerian Society of Experimental Biology*, 16 (2) 106-111.
- [7] **Carretto CFP, Almeida RBA, Furlan MR, Jorge AOC and Junqueira JC. 2010.** Antimicrobial activity of *Mentha piperita* L. against *Candida* spp. *Brazilian Dental Journal*, 13(1), 49.
- [8] **Diedhiou PM, Diop SAG, Mbaye N, Diedhiou I, Diallo Y, Djiba S, Faye R, Samb P. I. 2014.** Mango rotting in Southern Senegal, a big phytosanitary challenge. *International Journal of Biosciences*. 5(5):183-188.
- [9] **Doumbouya M, Abo K, Lepengue AN, Camara B, Kanko K, Aidara D, Koné D. 2012.** Activités comparées in vitro de deux fongicides de synthèse et de deux huiles essentielles, sur des champignons telluriques des cultures maraîchères en Côte d'Ivoire. *Journal of Applied Sciences* 50: 3520-3532.
- [10] **Edris AE, Farrag ES. 2003.** Antifungal activity of peppermint and sweet basil essential oils and their major aroma constituents on some plant pathogenic fungi from the vapor phase. *Nahr.* 47, 117-121.

- [11] **Furnery PM, Paolino D, Saija A, Marino A, Bisignano G. 2006.** *In vitro* antimycoplasmal activity of *Melaleuca alternifolia* essential oil. *Journal of Antimicrobial Chemotherapy*, 58: 706-707.
- [12] **Gautam AK, Kant M, Thakur Y. 2013.** Isolation of endophytic fungi from *Cannabis sativa* and study their antifungal potential. *Arch Phytopathol PI Prot*, 46: 627-635.
- [13] **Gil L, Tadesse W, Tolosana E, Lopez R. 2010.** *Eucalyptus* species management, history, status and trends in Ethiopia. Addis Abeba, Ethiopia Institute of Agricultural Research.
- [14] **Ghosch PG, Mukhopadhyay R, Gupta K. 2005.** Antifungal activity of the crude extracts and extracted phenols from gametophytes and sporophytes of two species of *Adiantum Taiwanianum*, 50(4):272-283.
- [15] **Govindarajin M, Sivakumar R, Rajeswary M, Yogalakshmi K. 2013.** Chemical composition and larvicidal activity of essential oil from *Ocimum basilicum* (L.) against *Culex tritaeniorhynchus*, *Aedes albopictus* and *Anophele subpictus* (Diptera: Culicidae). *Exp. Parasitol.* 134, 7-11.
- [16] **Lambert, R. J. W., Proteus N. Skandamis, Proteus J. Coote, et G.-JE Nychas. 2001.** « A study of the minimum inhibitory concentration and mode of action of oregano essential oil, thymol and carvacrol ». *Journal of applied microbiology* 91 (3): 453-462.
- [17] **Mahboubi M, Kazempour N. 2014.** Chemical composition and antimicrobial activity of peppermint (*Mentha piperita* L.) Essential oil, 36(1), 83-87.
- [18] **Matiz G, Osorio MR, Camacho F, Atencia M, Herazo J. 2012.** Effectiveness of antimicrobial formulations for acne based on orange (*Citrus sinensis*) and sweet basil (*Ocimum basilicum* L) essential oils. *Biomed. Rev. Inst. Nac. Salud*, 32, 125-133.
- [19] **Mbaye N. 2006.** Inventaire et caractérisation des champignons phytopathogènes responsables de maladies post-récolte des mangues (*Mangifera indica* L.) produites dans la zone des Niayes du Sénégal : cas des variétés Kent et Keitt, destinées à l'exportation. Doctorate Thesis Université Cheikh Anta Diop, Département de Biologie Végétale, Fac. Sciences, 121 p.
- [20] **Mokaddem H. 2011.** Etude phytochimique et biologique des espèces: *Eucalyptus globulus* (Myrtaceae), *Smyrniololus olusatrum* (Apiaceae), *Asteriscus maritimus* et *Chrysanthemum trifurcatum* (Asteraceae). Thèse de Doctorat. Option : Biochimie appliquée. Université Badji-Mokhtar, Annaba.
- [21] **Padman M, Janardhana GR. 2012.** Screening for inhibitory activities of essential oils on the growth of *Colletotrichum gloeosporioides* (Penz.) Penz. & Sacc., the causal agent of

leaf spot disease of *Murraya koenigii*., *Archives of Phytopathology and Plant Protection*, 45: 13, 1575-1581.

[22] **Pandey A, Yadava LP, Manoharan M, Chauhan UK, Pandey BK. 2012.** Effectiveness of cultural parameters on the growth and sporulation of *Colletotrichum gloeosporioides* causing anthracnose disease of Mango (*Mangifera indica* L.). *J Biol Sc*, 12: 123-13.

[23] **Pardo-De la Hoz CJ, Calderon C, Rincon AM, Cardenas M, Danies G, Lopez-Kleine L, Restrepo S and P. Jimenez P. 2016.** Species from the *Colletotrichum acutatum*, *Colletotrichum boninense* and *Colletotrichum gloeosporioides* species complexes associated with tree tomato and mango crops in Colombia. *Plant Pathology*, 65, 227-237.

[24] **Rao BPR, Kotharia SK, Rajput DK, Patel RP, Darokar MP. 2011.** Chemical and biological diversity in fourteen selections of four *Ocimum* species. *Nat. Prod. Commun.* 6, 1705-1710.

[25] **Shahwar D, Raza MA, Bukhari S, Bukkhari G. 2012.** Ferric reducing antioxidant power of essential oils extracted from *Eucalyptus* and *Curcuma* species. *Asian Pac J Trop Biomed*; 2 (Suppl 3): S1633-S1636.

[26] **Sokovic M, Glamoclija J, Marin PD, Brkic D, van Griensven LJLD. 2010.** Antibacterial effects of the essential oils of commonly consumed medicinal herbs using an in vitro model. *Mol. Basel Switz.* 15, 7532-7546.

[27] **Trevisan MTS, Vasconcelos Silva MG, Pfundstein B, Spiegelhalder B, Owen RW. 2006.** Characterization of the volatile pattern and antioxidant capacity of essential oils from different species of the genus *Ocimum*. *J. Agric. Food Chem.* 54, 4378-4382

[28] **Tsuchiya H, Sato M, Miyazaki T, Fujiwara S, Tanigaki S, Ohyama M, Tanaka T, Linuma M. 1996.** Comparative study on the antibacterial activity of phytochemical flavanones against methicillin-resistant *Staphylococcus aureus*. *J. Ethnopharmacol.* 50: 27-34.

[29] **Yoshimura H, Sawai Y, Tamotsu S, Sakai A, 2010.** 1,8- cineole inhibits both proliferation and elongation of BY-2 cultured tobacco cells, *J Chem Ecol*, 37(3):320-328.

[30] **Zani FG, Massimo S, Bianchi A, Albasini M, Melegari G, Vampa A, Belloti P, Mazza. 1991.** Studies on the genotoxic properties of essential oils with *Bacillus subtilis* rec-assay and Salmonella microsome reversion assay. *Planta Med.* 57:327-241.