

STUDY ON EFFECT OF DIFFERENT FEEDING REGIME IN CROSSBRED COWS DURING PRE-PARTUM PERIOD ON BIRTH WEIGHT OF CALVES

M.M. Islam* and S.V. Shah

Department of Livestock Production Management
College of Veterinary Science & Animal Husbandry, Anand Agricultural University
Anand-388001, Gujarat (India)

*E-mail: drislam37@gmail.com

Abstract: Eighteen crossbred cows were randomly selected and distributed in to three treatment groups comprising of 6 animals in each and fed with different feeding regimes during pre-partum period to study the effect of feeding on birth weight of calves. Average birth weight (kg) of calves during the experiment was maximum in T₃ (32.83±1.42) followed by T₂ (32.67±2.70) and T₁ (31.83±3.08) groups. However, there was no significant difference in birth weight of calves among treatment groups. Present study clearly indicated that the level of feeding during pre-parturient phase (last 45 days) had no effect on birth weight of calves.

Keywords: Feeding Regimes, Crossbred Cows, Birth Weight, Calves

Introduction

In developing countries like India and other third world countries, 70% of expenditure in dairy farming is on feeding of animals (Singh *et al.*, 2003). Either due to lack of awareness or due to unavailability of required quantity of feed, animals are under fed in field conditions (Khan *et al.*, 2004). Improper feeding of cows during pre-partum period could lead to low birth weight of newborn calf (Sasser *et al.*, 1988). Positive effect of steaming-up on birth weight of calves was reported by Das *et al.* (2007).

Materials and Methods

Present experiment was conducted on 18 HF x Kankrej crossbred cows which were randomly selected and distributed into three treatment groups comprising of 6 animals in each group. Animals were fed with different feeding regime during pre-partum period. Concentrate feed was not given to animals of T₁ (Farmer's feeding) group, given fixed @ 1kg/animal/d in T₂ (Modified feeding) group while in T₃ (Farm feeding) concentrate feed was given @ 500 g/animal/d in first week of experiment which was increased by 500 g every week till reaching to 3.5 to 4.0 kg/animal/d. Mineral mixture was not given to animals of T₁ group while fed @ 30 and 50 g/animal/d in T₂ and T₃ groups, respectively. Dry fodder was fed on *ad lib.* basis in

T₁ (paddy straw), T₂ (mixture of paddy and pigeon pea straw 50:50 ratio) and in T₃ (jowar hay). Daily 10 kg green hybrid napier (Co-3 variety) was fed to all animals irrespective of treatments. Birth weight of each calf was measured using electronic platform balance immediately after calving. Observations recorded during experimental period were statistically analyzed by one-way ANOVA using SPSS software 20.00 version.

Results and Discussion

Average fortnightly DMI (kg/100kg b.wt.) was significantly ($p < 0.05$) more in T₂ (1.78 ± 0.11) and T₃ (2.03 ± 0.15) as compared to T₁ (1.28 ± 0.08) group. Average DCPI (g/100kg b.wt.) was 52.08 ± 3.46 , 93.81 ± 4.28 and 129.88 ± 8.77 in T₁, T₂ and T₃ groups, respectively which differed significantly ($p < 0.05$) among each other. Average fortnightly TDNI (kg/100kg b.wt.) was significantly ($p < 0.05$) more in T₃ (1.17 ± 0.09) followed by T₂ (1.01 ± 0.07) when compared with T₁ (0.63 ± 0.04) group.

Average birth weight (kg) of calves during the experiment was maximum in T₃ (32.83 ± 1.42) followed by T₂ (32.67 ± 2.70) and T₁ (31.83 ± 3.08) group. However, there was no significant difference in birth weight of calves among treatment groups. Present study clearly indicated that the level of feeding during pre-parturient phase had no effect on birth weight of calves. In present study, up to 45 d before calving, animals of all treatment groups were maintained on same farm feeding. After start of experiment, it might have taken few days to alter/mobilize body reserve of animals. Experimental animals had a very short period of poor body reserve before calving which might not have influenced birth weight of calves. The absence of an effect of level of pre-partum feeding on calf birth weight is similar to the results of Prasad and Tomar (1996), where they had observed that animals kept on different feeding level in cows i.e. 80, 100 and 120 % of NRC ration delivered calves with body weight of 30.5, 31.5 and 32.87 kg, respectively. Panigrahi *et al.* (2005) also reported a non-significant effect of pre-partum feeding on birth weight of crossbred calves. Khan *et al.* (2004) reported significantly ($p < 0.05$) higher birth weight (kg) of calves from animals fed high energy (20.73 ± 2.24) as compared to low energy diets (17.87 ± 1.78) which is contradictory to the present study. Bindal (2012) also observed that calves of challenge fed group were 2.4 kg heavier than calves born to cows of control group but statistically the difference was not significant. Das *et al.* (2007) reported a positive effect of steaming-up on birth weight of calves. The mean birth weight of calves delivered by cows in steamed-up group was higher than the calves delivered by cows in control group (23.83 v/s 21.83 kg).

Conclusion

It may be concluded from the present study that level of feeding during pre-parturient phase (last 45 days) had no effect on birth weight of calves.

References

- [1] Bindal, Sameer. 2012. Effect of challenge feeding on performance and economics of milk production of crossbred cows. (M.V.Sc. thesis, National Dairy Research Institute, Karnal, Haryana, India).
- [2] Das, K.S., Das, N. and Ravikumar, G.V.P.P. 2007. Effect of 'steaming-up' on subsequent production in dairy cows. *Indian Journal of Animal Sciences*, **77(7)**: 583-585.
- [3] Khan, M.A.A., Islam, M.N., Khan, M.A.S. and Akbar, M.A. 2004. Effects of feeding high and low energy levels during late pregnancy on performance of crossbred dairy cows and their calves. *Asian-Aust. Journal of Animal Science*, **17(7)**: 947-953.
- [4] Panigrahi, B., Pandey, H.N. and Patnaik, A.K. 2005. Effect of prepartum feeding of crossbred cows on growth performance, metabolic profile and immune status of calves. *Asian-Australasian Journal of Animal Sciences*, **18**: 661-666.
- [5] Prasad, S. and Tomar, O.S. 1996. Effect of body condition at calving and postpartum level of feeding on the performance traits of crossbred cattle-body condition score and body weight. *Indian Journal of Dairy Science*, **49(2)**: 70.
- [6] Sasser, R.G., Williams, R.J., Bull, R.C., Ruder C.A. and Falk, D.G. 1988. Post-partum reproductive performance in crude protein restricted beef cows: return to estrus and conception. *Journal of Animal Sciences*, **66(12)**: 3033-3039.
- [7] Singh, J., Singh, B. M., Wadhwa, M. and Bakshi, M. P. S. 2003. Effect of level of feeding on the performance of crossbred cows during pre and post-partum periods. *Asian-Australasian Journal of Animal Sciences*, **16**: 1749-1754.